

Center-of-mass energy calibration in BES-III & VEPP-2000 experiments

Nickolai Muchnoi




Budker INP



February 24, 2020. Novosibirsk, Russian Federation

Laser backscattering for beam energy calibration

Taiwan Light Source: 1996
CO₂ laser & HPGe detector



$E = 1305.8 \pm 1.7 \text{ MeV}$
Phys. Rev. E v.54 (5) 1996

STORAGE RINGS

- ▶ BESSY-I - 1998
- ▶ BESSY-II - 2002
- ▶ VEPP-3 - 2008
- ▶ NewSUBARU - 2009
- ▶ ANKA - 2015


e^+e^- COLLIDERS

- ▶ VEPP-4M - 2005
- ▶ BEPC-II - 2010
- ▶ VEPP-2000 - 2012

Inverse Compton Scattering

When $\gamma \gg 1$:

$\varepsilon_0, \varepsilon, \omega \gg \omega_0$



$$\omega_0^* = \frac{hc}{\lambda_0} \cos^2(\alpha/2)$$

Scattering energies:

$$\max(\omega) = \kappa \cdot \varepsilon_0 / (1 + \kappa)$$

$$\min(\varepsilon) = \varepsilon_0 / (1 + \kappa)$$

Universal scattering parameter:

$$u = \frac{\omega}{\varepsilon} = \frac{\theta_\varepsilon}{\theta_\omega} = \frac{\omega}{\varepsilon_0 - \omega} = \frac{\varepsilon_0 - \varepsilon}{\varepsilon}$$

is in the range


$$0 \leq u \leq \kappa, \text{ where } \kappa = \frac{4\omega_0^* \varepsilon_0}{(mc^2)^2}.$$

$$\kappa \simeq 0.0153 \text{ for } \varepsilon_0 = 1 \text{ GeV and } \omega_0^* = 1 \text{ eV}$$

Scattering angles: $\gamma\theta_\omega = \sqrt{\kappa/u - 1}$; $\theta_\varepsilon = u\theta_\omega$


$\max(\theta_\varepsilon) = 2\omega_0^*/mc^2$ ($\simeq 10$ urad for green light).

BEPC-II Collider




- ▶ SR losses: $E_{ip} = E_{BEMS} + 4.75 \cdot 10^{-6} (E_{BEMS})^4$ [GeV]
- ▶ IP angle: $E_{cm} = 2 \cos(0.011) \sqrt{E_{ip}^{e^+} E_{ip}^{e^-}}$

Beam Energy Measurement System




γ -rays detection



γ -rays spectrum example

electron: 2018.04.22 [23:43:18 - 06:28:50] 2018.04.23. Live-time: 3 hours 16 min 16 s (10 files).




Absolute Scale Calibration

Source	γ -rays energies, keV	Reference *
^{137}Cs	661.657 ± 0.003	vol.4, 2008
^{60}Co	1173.228 ± 0.003 1332.492 ± 0.004	vol.4, 2008
^{228}Ac (^{232}Th)	911.209 ± 0.006	vol.6, 2011
^{212}Bi (^{232}Th)	727.330 ± 0.030 1620.740 ± 0.010	vol.2, 2004
^{208}Tl (^{232}Th)	583.187 ± 0.002 860.560 ± 0.030 2614.511 ± 0.010	vol.2, 2004

* Table of Radionuclides, Bureau International des Poids et Mesures
<https://www.bipm.org/en/publications/scientific-output/monographie-ri-5.html>

Photopeak Model



Photopeak Model


$$f(x) = B + \frac{C}{2} \operatorname{erfc} \left(\frac{x}{\sqrt{2}\sigma_R} \right) + \frac{N}{N_1} \begin{cases} \exp \left(-\frac{x^2}{2\sigma_R^2} \right) & \text{if } x > 0; \\ \exp \left(-\frac{x^2}{2\sigma_L^2} \right) & \text{if } -\kappa\sigma_L < x \leq 0; \\ \exp \left(\frac{\kappa x}{\sigma_L} + \frac{\kappa^2}{2} \right) & \text{if } x \leq -\kappa\sigma_L \end{cases}$$

Here $x = (E - E_{\max})$ is the difference between the energy deposition in the detector and its most probable value, B is the background level, N is the number of counts in the photopeak while N_1 is the normalization constant:

$$N_1 = \sqrt{\frac{\pi}{2}}\sigma_R + \left\{ \frac{1}{\kappa} \exp \left(-\frac{\kappa^2}{2} \right) + \sqrt{\frac{\pi}{2}} \operatorname{erf} \left(\frac{\kappa}{\sqrt{2}} \right) \right\} \sigma_L$$


Photopeak Fit Example

^{60}Co (1332.492 keV) $\chi^2/\text{ndf} = 13.0/14$




Photopeak Fit Example


^{208}Tl (2614.511 keV) $\chi^2/\text{ndf} = 21.7/20$



Energy Resolution




Absolute Energy Scale



Compton Edge in γ -spectrum




positron: 2018.04.27 [19:20:24 - 12:31:37] 2018.04.28. Live-time: 4 hours 21 min 5 s (16 files).




$$E_{\text{beam}} = \frac{\omega_{\text{max}}}{2} \left(1 + \sqrt{1 + \frac{m^2}{\omega_0 \omega_{\text{max}}}} \right), \quad \omega_0 = 0.11706523 \text{ eV}.$$

BEMS check at J/ψ - 2018




BEMS check at $\psi(2S)$ - 2018




Phys. Rev. D 90, 012001 (2014)



"Precision measurement of the mass of the τ lepton"



VEPP-2000 e^+e^- collider



Circumference:

$$P = 24.4 \text{ m}$$

C. m. s. energy:


$$E = 200 - 2000 \text{ MeV}$$

Luminosity / i.p.:

$$L = 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$$


Could not find a free straight section for laser i.p.!

VEPP-2000 BEMS




Spectrum $\varepsilon_0 = 640 \text{ MeV}$, $\lambda_0 = 5.426463 \text{ }\mu\text{m}$

2019.06.24 [18:35:21 - 19:15:28] 2019.06.24. Live-time: 0 hours 35 min 32 s (2 files).



Interference



$\theta_{int} \gg \theta_{rad}$: take only $\phi = 0$

$$\text{Time for electron } A \rightarrow B \rightarrow C: t_e = \frac{2R\theta}{\beta c}$$

$$\text{Time for photon } A \rightarrow C: t_\gamma = \frac{2R \sin \theta}{c} \cos \psi$$

$$\text{Phase advance: } \Delta\Phi = 2\pi c \left[\frac{t_e}{\lambda} - \frac{t_\gamma}{\lambda} - \frac{2t_e}{\lambda_0} \right]$$

λ_0 - laser wavelength. Since $\theta, \psi, 1/\gamma \ll 1$

$$\Delta\Phi \simeq \frac{\omega R}{c} \left[\theta \left(\frac{1}{\gamma^2} - \frac{4\omega_0}{\omega} + \psi^2 \right) + \frac{\theta^3}{3} \right].$$

For 1 MeV photon $\lambda = 1.24 \cdot 10^{-12}$ m. For $R = 140$ cm, $E = 1$ GeV,

$\Delta\Phi = 2\pi$ when $\theta \simeq 0.1/\gamma$ and $\overline{AC} \simeq 0.1$ mm $\simeq 10^8 \lambda$!

Energy spectrum

$$\frac{d\dot{N}_\gamma}{d\omega d\psi} \propto \omega^{1/3} \text{Ai}^2(x), \text{ where}$$



$$x = \left[\frac{\omega R}{2\hbar c} \right]^{2/3} \left[\frac{1}{\gamma^2} - \frac{4\omega_0}{\omega} + \psi^2 \right].$$

$$\frac{d\sigma}{d\omega} = \frac{d\sigma_{KN}}{d\omega} \int_z^\infty \text{Ai}(x) dx, \text{ where}$$

$$z = (u/\chi)^{2/3} (1 - \kappa/u), \text{ where}$$

$$\chi = \gamma \frac{B}{B_c}, \quad u = \frac{\omega}{\varepsilon_0 - \omega}, \quad \kappa = 4 \frac{\varepsilon_0 \omega_0}{m^2 c^4}$$

$$B_c = \frac{m^2 c^2}{\hbar e} = 4.414 \cdot 10^9 \text{ [T]}$$




$$\omega_0 = 0.117 \text{ eV}; \quad \varepsilon_0 = 900 \text{ MeV}; \quad R = 140 \text{ cm}$$


$$\text{Ai}(x) = \frac{1}{\pi} \int_0^\infty \cos\left(xt + \frac{t^3}{3}\right) dt$$

The Fit Result

2019.06.24 [18:35:21 - 19:15:28] 2019.06.24. Live-time: 0 hours 35 min 32 s (2 files).



$e^+e^- \rightarrow K_S K_L$, CMD-3 (2013)



Summary

- ▶ Laser back-scattering is being applied now for c. m. s. energy calibration on two lepton colliders - BEPC-II and VEPP-2000.
- ▶ Center-of-mass energy was calibrated with accuracy $\gtrsim 30 \text{ keV}$ in the energy range from 360 MeV to 3700 MeV.
- ▶ The new τ -lepton mass measurement was performed in 2018 by BES-III detector. The new result is expected to arrive this year.

Thank You!